

One-dimensional dynamics for traveling fronts in coupled map lattices

R. Carretero-González,* D. K. Arrowsmith, and F. Vivaldi

School of Mathematical Sciences, Queen Mary and Westfield College, Mile End Road, London E1 4NS, United Kingdom

(Received 1 April 1999)

Multistable coupled map lattices typically support traveling fronts, separating two adjacent stable phases. We show how the existence of an invariant function describing the front profile allows a reduction of the infinitely dimensional dynamics to a one-dimensional circle homeomorphism, whose rotation number gives the propagation velocity. The mode locking of the velocity with respect to the system parameters then typically follows. We study the behavior of fronts near the boundary of parametric stability, and we explain how the mode locking tends to disappear as we approach the continuum limit of an infinite density of sites.

PACS number(s): 05.45.Ra

I. INTRODUCTION

Coupled map lattices (CML's) are arrays of low-dimensional dynamical systems with discrete time, originally introduced in 1984 as simple models for spatiotemporal complexity [1]. CML's have been extensively used in modeling spatiotemporal chaos in fluid phenomena such as turbulence [2], convection [3], and open flows [4]. Equally important is the analysis of pattern dynamics, which has found applications in chemistry [5] and patch population dynamics [6]. One important feature of pattern dynamics is the existence of traveling fronts, which occur at the pattern boundaries, and are also seen to emerge from apparently decorrelated media [7]. This paper extends the work on the behavior of a traveling interface on a lattice developed in [8–11]. Our main results are (i) a constructive procedure for the reduction of the infinitely dimensional dynamics of a front to one dimension; (ii) a characterization of the behavior of fronts near the boundary of parametric stability; (iii) a characterization of the behavior of fronts near the continuum limit.

We consider a one-dimensional infinite array of sites. At the i th site there is a real dynamical variable $x(i)$, and a local dynamical system—the *local map*. The latter is given by a real function f which we assume to be the same at all sites. The dynamics of the CML is a combination of local dynamics and coupling, which consists of a weighted sum over some neighborhood. The time evolution of the i th variable is given by

$$x_{t+1}(i) = \sum_k \varepsilon_k f(x_t(i+k)),$$

where the range of summation defines the neighborhood. The coupling parameters ε_k are site independent, and they satisfy the conservation law $\sum \varepsilon_k = 1$, to prevent unboundedness as time increases to infinity. The two most common choices for the coupling are

$$x_{t+1}(i) = (1 - \varepsilon)f(x_t(i)) + \varepsilon f(x_t(i-1)) \quad (1)$$

and

$$x_{t+1}(i) = (1 - \varepsilon)f(x_t(i)) + \frac{\varepsilon}{2}[f(x_t(i-1)) + f(x_t(i+1))], \quad (2)$$

which are called *one-way* and *diffusive* CML's, respectively. The diffusive CML corresponds to the discrete analogue of the reaction-diffusion equation with a symmetrical neighboring interaction. There is now a single coupling parameter ε which is constrained by the inequality $0 \leq \varepsilon \leq 1$, to ensure that the sign of the coupling coefficients in Eqs. (2) and (1) (i.e., ε , $\varepsilon/2$, and $1 - \varepsilon$) remains positive.

In this paper we study front propagation in *bistable* CML's. The local mapping f is continuous and has two stable equilibria, and a *front* is any monotonic arrangement of the state variables, linking the two equilibria asymptotically.

We will show how to construct a one-dimensional circle map describing the motion of the front. Such a mapping originates from the existence of an invariant function describing the asymptotic front profile, and of a one-dimensional manifold supporting the transient motions. The rotation number of the circle map will then give the velocity of propagation, resulting in the occurrence of *mode locking*, i.e., the parametric stability of the configurations that correspond to *rational* velocity. We will describe the vanishing of this phenomenon in the continuum limit, as the width of the front becomes infinite. We shall also be concerned with the evolution of the front shape near the boundary of parametric stability, where the continuity of the local map ensures a smooth evolution of the front shape.

Velocity mode locking is commonplace in nonlinear coupled systems (e.g., Frenkel-Kontorova models [12], Josephson-junction arrays [13], excitable chemical reactions [14], and nonlinear oscillators [15]); the present work provides further support for its genericity, and highlights key dynamical aspects.

Throughout this paper, the very existence of fronts in the regimes of interest to us is inferred from extensive numerical evidence. We are not concerned with existence proofs here. Fronts have been proved to exist in various situations,

*Present address: Department of Mathematics and Statistics, Simon Fraser University, Burnaby, BC, Canada V5A 1S6. Electronic address: ricardo_carretero@sfu.ca, URL: <http://www.math.sfu.ca/~rcarretero/ric.html>

mainly for *discontinuous* piecewise affine maps (see [11] and references therein); in the present context, however, continuity is crucial.

Following [9], we consider a CML whose local map f is continuous and monotonically increasing and which possesses exactly two stable fixed points x_-^* and x_+^* . It then follows that there exists a unique unstable fixed point x^* such that $x_-^* < x^* < x_+^*$. The homogeneous fixed states $x(i) = x_\pm^*$, $\forall i \in \mathbb{Z}$, inherit the stability of the fixed points x_\pm^* [16]. We denote by $I_- = [x_-^*, x^*)$ and $I_+ = (x^*, x_+^*]$ the basins of attraction of x_-^* and x_+^* , respectively, while $I = [x_-^*, x_+^*]$.

A *minimal mass state* is a state satisfying the monotonicity condition $x(i) \leq x(i+1)$, for all i . It can be shown directly from the system equation that the image of a minimal mass state has the same property. A *front* is a minimal mass state satisfying the asymptotic condition $\lim_{i \rightarrow \pm\infty} x(i) = x_\pm^*$. The main properties of a front are its *center of mass* μ_t and its *width* σ_t^2 , which measure its position and spread at time t , respectively. They are defined as the mean and variance of the variable i with respect to the time-dependent probability distribution

$$p_t(i) = \frac{|\Delta x_t(i)|}{\sum_{i=-\infty}^{\infty} |\Delta x_t(i)|}, \quad (3)$$

where $\Delta x_t(i) = x_t(i+1) - x_t(i)$ is the variation of the local states. We have

$$\mu_t = \sum_{i=-\infty}^{\infty} i p_t(i), \quad (4)$$

$$\sigma_t^2 = \sum_{i=-\infty}^{\infty} (i - \mu_t)^2 p_t(i).$$

A state $X_t = \{x_t(i)\}$ with finite center of mass and width is said to be *localized*.

In this paper we are interested in fronts of *fixed shape*, moving at velocity v . They are described by the equation

$$x_t(i) = h(i - vt), \quad v = \lim_{t \rightarrow \infty} \frac{\mu_t}{t}, \quad t, i \in \mathbb{Z}. \quad (5)$$

Here the function $h: \mathbb{R} \rightarrow [x_-^*, x_+^*] = I$ is to be determined subject to the condition that it be monotonic, with $\lim_{x \rightarrow \pm\infty} = \pm x_\pm^*$. The degree of smoothness of h will depend on the regime being considered.

The object of interest to us is the central part of the front. Far away from the center, the lattice is almost homogeneous [i.e., $|\Delta x_t(i)| \ll |I|$], and the dynamics is dominated by the attraction toward the stable points of the local map. The qualitative evolution of the center of the front can be understood as the result of the competition between local dynamics and coupling (see Fig. 1, for the one-way case). For small ε , the attraction toward the fixed points x_\pm^* overcomes the effect of the coupling, resulting in propagation failure (zero velocity) [9]. A sufficiently large coupling will instead cause a site located within the basin I_+ to switch to the basin I_- ,




FIG. 1. The dynamics of a front for a one-way CML results from the competition between local dynamics and coupling. The schematic contributions from the local dynamics (arrows with filled arrowhead) and coupling (arrows with empty arrowhead) are depicted for all front sites at time t (filled circles). A sufficiently large coupling causes a site located within the basin I_+ (filled circle at the center of front) to switch to the basin I_- , and move rapidly toward x_-^* . As a consequence, the center of mass of the front will move to the right, resulting in propagation.

and move rapidly toward x_-^* . As a consequence, the center of mass of the front will move to the right, resulting in propagation.

A similar argument can be applied in the diffusive case. Now however the coupling is symmetric, and a bias to either of the stable points will have to be introduced via an asymmetry in the local map. For instance, increasing the size of the basin of attraction of x_-^* will result in propagation from left to right for an increasing front.

In previous works we have shown that the dynamics of a *finite-size* interface in a class of piecewise linear one-way CML's can be reduced to a single one-dimensional map [9,10]. The finiteness of the front depended on the existence of degenerate superstable fixed points of the local map, which caused nearby orbits to collapse onto the stable states in a single iteration. In this paper we remove such degeneracy, and consider smooth local maps and infinitely extended fronts (the case of a discontinuous local map was treated in [11]). We shall provide evidence that every front evolves toward a unique asymptotic regime, characterized by a constant velocity as well as an invariant shape. Under these assumptions, we then show how the front behaves at the boundary of the regions of parametric stability (here the continuity of the local map is essential), and how the reduction to one-dimensional dynamics can be achieved.

This paper is organized as follows. In Sec. II we describe the behavior of traveling fronts in the continuum limit, when the density of interfacial sites is large. We obtain an ordinary differential equation (ODE) describing the shape of the traveling front, and with it we find new classes of fronts. In Sec. III we consider the asymptotic shape of the front, and we provide extensive evidence that such a shape is fixed and is described by a continuous function. This result allows us to derive a procedure for the reduction of the infinite-dimensional interface dynamics to a one-dimensional problem described by the *auxiliary map*. In Sec. IV we show that the auxiliary map is a circle map and we relate its rotation number to the velocity of the front, from which the mode locking of the velocity with respect to the system parameters follows. Finally, we explain in terms of reduced dynamics the vanishing effect of mode locking when the continuum limit is approached.

II. THE CONTINUUM LIMIT

In this section we consider fronts with large widths, for which the relative density of sites is large, and the continuum approximation becomes appropriate. To achieve a front with such features, the attraction toward x_{\pm}^* and the repulsion of x^* must be small. Because f is continuous and monotonic, then f is necessarily close to the identity, i.e.,

$$\delta_f = \sup_{x_-^* < x < x_+^*} |f(x) - x| \ll 1.$$

Choosing functions f such that $\delta_f \rightarrow 0$ is referred to as the *continuum limit*.

Inserting Eq. (5) into the equations of motion (1) and (2) we find that

$$(a) \quad h(z - v) = (1 - \varepsilon)f(h(z)) + \varepsilon f(h(z - 1)), \quad (6)$$

$$(b) \quad h(z - v) = (1 - \varepsilon)f(h(z)) + \frac{\varepsilon}{2}[f(h(z - 1)) + f(h(z + 1))],$$

for the one-way and diffusive CML's, respectively, where $z = i - vt$. A function h satisfying the functional equation (6) represents the fixed shape of a front traveling at the velocity v .

To solve Eq. (6) in the continuum limit, we assume f and h to be twice differentiable, and consider the Taylor series of h in z , up to second order. The Taylor expansion becomes accurate as the width increases, since in this case the variation of h over adjacent lattice sites tends to zero. We obtain

$$h(z) - f(h(z)) + Ah'(z) - \left(\frac{\varepsilon f''(h(z))}{2}\right)h'(z)^2 + \left(\frac{v^2 - \varepsilon f'(h(z))}{2}\right)h''(z) = 0, \quad (7)$$

where $A = [\varepsilon f'(h(z)) - v]$ and $A = -v$, for the one-way and diffusive CML's, respectively. In the continuum limit we can further simplify Eq. (7) by considering $f'(x) = 1$ and $f''(x) = 0$, to obtain

$$(a) \quad h(z) - f(h(z)) + \left(\frac{\varepsilon(\varepsilon - 1)}{2}\right)h''(z) = 0, \quad (8)$$

$$(b) \quad h(z) - f(h(z)) - vh'(z) + \left(\frac{v^2 - \varepsilon}{2}\right)h''(z) = 0,$$

for the one-way and diffusive CML's, respectively, where we set $v = \varepsilon$ in the one-way case since in the continuum limit $f(x) \rightarrow x$ and thus the rate of information exchange (i.e., the velocity) is equal to ε . For the diffusive case the velocity is not equal to ε , since the total information exchange comes from the competition between the left and right neighbors. Nevertheless, as we shall see, it is possible to give an analytical approximation to the velocity for the case of an asymmetric cubic local map.

Equations (8) are similar to those obtained in [17], where the traveling front in a lattice of coupled ODE's is reduced to




FIG. 2. Qualitative features of the phase space $h'(t)$ vs $h(t)$ of the ODE's (8), corresponding to the traveling front solution in the continuum limit. (a) A one-way CML corresponds to Hamiltonian motions. (b) A diffusive CML corresponds to dissipative motions. Note that in (b) a heteroclinic connection between unstable points can still exist in the presence of friction.

a single equation. The ODE's (8) describe the motion of a particle of mass $m = (v^2 - \varepsilon)/2$, subject to the potential $V(x) = \int [f(x) - x] dx$, with maxima located at the stable fixed points of the local map (Fig. 2).

In the one-way case, the system is conservative. For numerical experiments, we choose a symmetric local map f with fixed point $x_{\pm}^* = \pm 1$ and $x^* = 0$. The resulting potential is also symmetric. There exist two heteroclinic connections, joining x_-^* to x_+^* and x_+^* to x_-^* , respectively [the thick lines in Fig. 2(a)]. They correspond, respectively, to an increasing and a decreasing symmetric traveling front for the CML.

In the diffusive case, the differential equation has the dissipative term $-vh'(z)$. For the local map, we choose $0 < x^* = p < 1$, which introduces an asymmetry in the system, and the maxima of the potential are now unequal: $V(x_-^*) > V(x_+^*)$. Imposing a heteroclinic connection from x_-^* and x_+^* constrains the velocity v of the front (see below). For larger velocities, the separatrix emanating from x_-^* approaches p , while for smaller v it escapes to infinity. Since the presence of friction breaks the time-reversal symmetry, only one heteroclinic connection is possible, and the separatrix emanating from x_+^* always approaches p [the thick lines in Fig. 2(b)].

The continuum approximation can be used to construct new kinds of traveling fronts. For example, the librating orbits in Fig. 2(a) (one-way case), correspond to spatially periodic traveling fronts that never touch the stable points [see Fig. 3(a) (iii)]. Such spatially periodic orbits do not exist in the diffusive case. Nevertheless, it is possible to construct the traveling front departing from x_+^* that dissipates down to p . This new solution has a damped oscillatory profile [see Fig. 3(b) (i)].

In the remainder of this section, we briefly examine the case of a cubic local map, providing the dominant behavior of a general bistable local map in the continuum limit. We use the one-parameter families of cubics

$$(a) \quad f(x) = \frac{x}{2}[3 - v - (1 - v)x^2], \quad (9)$$

$$(b) \quad f(x) = (1 - v)(px^2 - x^3 - p) + (2 - v)x,$$




FIG. 3. Traveling front solutions in the continuum limit approach. (a) Heteroclinic [(i) and (ii)], and oscillatory (iii) solutions in a one-way CML. (b) Damped heteroclinic solutions: (i) connects the stable fixed point x_+^* to the unstable fixed point $x^*=p$; (ii) connects the two stable points.

for the one-way and diffusive CML's, respectively. Again, $x_{\pm}^* = \pm 1$ for both cases, while $x^*=0$ in the one-way case and $x^*=p$ in the diffusive case, where $0 < p < 1$ controls the asymmetry. The continuum limit is attained by letting the parameter ν approach 1 from below. Substituting the cubic local maps (9) in the differential equations (8), one finds expressions for the heteroclinic connections corresponding to the traveling front solutions:

$$\begin{aligned} \text{(a)} \quad h(z) &= \tanh\left(\sqrt{\frac{1-\nu}{2\varepsilon(1-\varepsilon)}}z\right), \\ \text{(b)} \quad h(z) &= \tanh\left(\frac{(1-\nu)p}{\nu}z\right), \end{aligned} \quad (10)$$

where

$$\begin{aligned} \text{(a)} \quad \nu &= \varepsilon, \quad \sigma^2 = \frac{2\pi^2}{3} \frac{\varepsilon(1-\varepsilon)}{1-\nu}, \\ \text{(b)} \quad \nu &= p\sqrt{\varepsilon(1-\nu)}, \quad \sigma^2 = \frac{\pi^2}{3} \frac{\varepsilon}{1-\nu}, \end{aligned} \quad (11)$$

for the one-way and diffusive CML's, respectively. In the diffusive case, the expression for the velocity is derived from imposing a heteroclinic connection, while the scaling of the width σ^2 is found from the solutions (10). Note that for both models the functional dependence of the width on the parameter ν is the same, and it describes the rate at which the front broadens as the continuum limit is approached. Moreover, from Eqs. (10) and (11) we have that in the diffusive case h is independent of p .

While in the continuum limit the front is described by a continuous function h [cf. Eq. (10)], there is no *a priori* reason why such a function should continue to exist away from the limit, due to the discrete nature of the system. We shall nonetheless give evidence that the dynamics of a front far from the continuous limit remains one dimensional.

III. REDUCED DYNAMICS OF THE TRAVELING FRONT

In this section we provide evidence that every front has a fixed profile, which can be characterized by an invariant function h . Such a function will then be used to construct a




FIG. 4. The traveling front shape is reconstructed by superimposing snapshots of the discrete interface in a comoving reference frame. (a) One-way coupling: $f(x) = \tanh(x/0.2)$, $\varepsilon = 0.398\,011$, $\nu(\varepsilon) \approx 0.285\,603\,1 \approx 2/7$. (b) Diffusive coupling: f is the second iterate of the logistic map, with $\varepsilon = 0.2$ and $\nu(\varepsilon) \approx 0.009\,791\,5$. (c) Diffusive coupling: f as in (b), with $\varepsilon = 0.6$ and $\nu(\varepsilon) \approx 0.111\,827\,3$.

one-dimensional mapping describing the front evolution—the auxiliary map.

If the velocity ν of the front is *irrational*, then the collection of points $i - \nu t$, with i and t integers, form a set dense on the real line. Numerical experiments consistently suggest that in the case of a front, the closure of the set of points $(i - \nu t, x_t(i)) \in \mathbb{R}^2$ forms the graph of a continuous and monotonic function: $h: \mathbb{Z} \rightarrow [x_-^*, x_+^*]$, which is a solution to the functional equation (6).

The results for both CML models are summarized in Fig. 4, where we have superposed all translates of the discrete fronts, after eliminating transient behavior. This procedure requires computing ν numerically, which was done using some 10^7 – 10^8 iterations of the CML. [In principle, a numerical solution to Eq. (6) can be found using various iterative functional schemes. However, all the schemes considered were plagued by slow convergence and are not discussed here.]

In the case in which $\nu = p/q$ is *rational*, the function h is specified only at a set of q equally spaced points. It turns out, however, that the definition of h becomes unequivocal in a prominent parametric regime, corresponding to the boundary of the so-called mode-locking region or *tongue*. The latter is defined as the collection of parameters (ε, ν) corresponding to a given rational velocity, where ν (not necessarily one dimensional) parametrizes the family of local maps—for the one-way CML we typically use $f(x) = \tanh(x/\nu)$.

We defer the discussion of the origin of such regions to the next section. Here we consider a sequence of parameters $(\varepsilon_n, \nu_n) \rightarrow (\varepsilon_*, \nu_*)$, converging from the outside toward a boundary point (ε_*, ν_*) of the tongue (see Fig. 5). Independently from the path chosen to approach the boundary point, the front h appears to approach a unique limiting shape. The limiting shape is a step function with q steps (where ν




FIG. 5. Approximating traveling front for a rational velocity. The parametric point located at the edge of a tongue (small box), is approached both transversally (path A) and tangentially (paths B and C). The tongue corresponds to a traveling front with velocity $v(\epsilon)=1/3$, which is periodic with period 3. In all cases the front shape approaches the same step function, with 3 steps per unit length in z . Note that the fronts have been shifted for clarity. Here the local map is $f(x)=\tanh(x/v)$, and the parameters at the boundary of the mode-locking region are $(\epsilon_*, \nu_*)=(0.3983, 0.1)$.

$=p/q$) for every unit length—the horizontal length of each step is $1/q$ since there are q equidistant points in every horizontal interval of unit length for a $v=p/q$ orbit. In the limit, the front dynamics becomes periodic, with periodic points corresponding to the midpoint of each step. This observation suggests that choosing step fronts with the periodic points at their midpoints ensures continuity of the front shapes at the resonance tongue boundaries.

In the next section, we shall explain this phenomenon in terms of the dynamics of a one-dimensional map—the auxiliary map Φ —which we now define. The idea is to describe the evolution of any site in the front by means of a single site, the *central site* $\bar{x}_i(0)$, defined as the site that is closest to the unstable point x^* . The position of the central site moves along the lattice with an average velocity $v(\epsilon)$, since it follows the center of the interface. Following [9,10], we define the map Φ as

$$\bar{x}_{i+1}(0) = \Phi(\bar{x}_i(0)). \tag{12}$$

If the velocity is irrational, the domain of definition of the map is a set of points dense in an interval (see next section), and the possibility exists of extending Φ continuously to the interval. In Fig. 6(a) and (b), we plot the graph of Φ for a one-way and a diffusive CML, respectively. The auxiliary map corresponds to the square region depicted with thick lines, while the other regions represent delay Poincaré maps of some neighboring sites. Indeed, for each neighbor j of the central site, there is a corresponding auxiliary circle map Φ_j , such that $\bar{x}_{i+1}(j) = \Phi_j(\bar{x}_i(j))$, with $\Phi = \Phi_0$ (see below).

If the velocity is rational, Eq. (12) defines Φ only at a finite set of points, and to extend the domain of definition,





FIG. 6. Auxiliary maps Φ for the central site of the interface defined in the square region depicted by the thick lines. (a) One-way CML: local map $f(x)=\tanh(x/0.2)$ with $\epsilon=0.4$, $\nu=0.28973453$. (b) Diffusive: same parameters as in Fig. 4(c). The delay Poincaré section $\Phi(x)$ corresponds to the central rectangular region of each plot (region 1) in (a). Each rectangular region corresponds to the return map for a particular combination of sites. For instance, region 2 in (a) corresponds to $\bar{x}_{i+1}(1)$ vs $\bar{x}_i(1)$.

one must make use of Eq. (12) on suitable transients. We have verified numerically that when a front is perturbed, the perturbation relaxes quickly onto a one-dimensional manifold, along which the original front is approached. The process of randomly disturbing the front amounts to a random walk path reconstruction of the one-dimensional manifold. Such one-dimensional transients were found to be independent of the detail of the perturbation, giving an unequivocal definition of the auxiliary map in the rational case also. This is illustrated in Fig. 7. Crucially, this construction yields a map that changes continuously within the tongue, matching the the behavior at the boundary of the tongue. Thus we




FIG. 7. Typical reconstruction of the auxiliary map Φ inside a mode-locking tongue. The large stars locate the original periodic orbit well inside a tongue (in this example $v=1/5$). A small random perturbation is periodically applied to the central site of the front. The state of each perturbed front is depicted by circles. After few transient iterations (2 or 3), the perturbed front relaxes onto the one-dimensional manifold represented by the thick line. This technique is applied repeatedly until the whole one-dimensional manifold is filled in.

conjecture that the auxiliary map Φ depends continuously on the coupling parameter ε . In the next section we shall explore some consequences of the continuity.

We finally relate the dynamics of the entire front to that of the central site, governed by $\Phi(x)$. Let $\bar{x}_i(j)$ denote the j th neighboring site of the central site $\bar{x}_i(0)$, where j is positive (negative) for the right (left) neighbors. The dynamics of $\bar{x}_i(j)$ can be deduced from that of $\bar{x}_i(0)$ and the knowledge of h , as follows:

$$\bar{x}_i(j) = h \circ \tau_j \circ h^{-1}(\bar{x}_i(0)) \quad (13)$$

where τ_j is the translation by j on \mathbb{R} . Since $\Phi_j(x)$ maps $\bar{x}_i(j)$ to $\bar{x}_{i+1}(j)$, the pair $(\bar{x}_i(j), \bar{x}_{i+1}(j))$ belongs to the graph of Φ_j . By applying the operator $h \circ \tau_j \circ h^{-1}$ to the function $\Phi(\bar{x}_i(0))$ we obtain

$$\begin{aligned} h \circ \tau_j \circ h^{-1} \Phi(\bar{x}_i(0)) &= h \circ \tau_j \circ h^{-1}(\bar{x}_{i+1}(0)) \\ &= \bar{x}_{i+1}(j) = \Phi_j(\bar{x}_i(j)), \end{aligned}$$

where we used Eq. (13) which relates neighboring sites. Thus $h \circ \tau_j \circ h^{-1}$ provides a conjugacy between Φ and Φ_j and enables us to reconstruct the whole interfacial dynamics from the behavior of the central site.

IV. MODE LOCKING OF THE PROPAGATION VELOCITY

In this section we show that the auxiliary map Φ is a circle homeomorphism (see Fig. 8). The mode locking of the




FIG. 8. The auxiliary map Φ , accounting for the dynamics of the site closest to the unstable point x^* , is a circle map on $[x^*-a, x^*+a]$, with two increasing branches f_+ and f_- .

front velocity will then follow from the mode locking of the rotation number of Φ . Furthermore, the conjectured continuous dependence of Φ on ε implies a continuous dependence of the rotation number on ε , and in particular, Φ takes all rotation numbers between any two realized values. For instance, the front velocity in a one-way CML takes the values 0 and 1 for $\varepsilon=0$ and 1, respectively, and thus as the coupling parameter varies, all velocities $v \in [0, 1]$ are realized. For a diffusive CML only an interval $[0, v_{max}]$ is attained since the maximum velocity $v_{max} = v(\varepsilon=1)$ does not reach 1 because of the competition between the attractors.

Let us consider a continuous and increasing traveling front $h(i-vt+i_0)$ with positive irrational velocity $0 < v < 1$. The largest possible separation between $\bar{x}_i(0)$ and x^* corresponds to the position of h for which two consecutive points on the lattice are equally spaced from the unstable point x^* . Suppose that the front shape h is positioned such that for site i , we have $h(i) = x^*$. We choose α such that

$$h(i-\alpha) = x^*-a \quad \text{and} \quad h(i+1-\alpha) = x^*+a \quad (14)$$

where $0 \leq \alpha \leq \min(|x_+^* - x^*|, |x_-^* - x^*|)$. By adding the two equations in (14) one obtains an equation for α , and a can then be evaluated. If the front is at a position where it satisfies the equations (14) for some i , then the i th and $(i+1)$ th sites are equally spaced from x^* , and the dynamics of the site closest to x^* is contained in the interval $[x^*-a, x^*+a]$. Any shift of the front will cause either one of the two sites to be closer to x^* than originally.

We now follow the dynamics of $\bar{x}_i(0)$ in $[x^*-a, x^*+a]$. Suppose that at time τ the i th site is the closest to x^* so $\bar{x}_\tau(0) = x_\tau(i)$. We want to know which site will be closest to x^* at time $\tau+1$. Since we are considering the case $v > 0$ there are two possibilities: (a) the i th site again $[\bar{x}_{\tau+1}(0) = x_{\tau+1}(i)]$ or (b) the $(i+1)$ th site $[\bar{x}_{\tau+1}(0) = x_{\tau+1}(i+1)]$. Redefining $h_i(i) = h(i-vt+i_0)$, we find two cases,

$$h_{\tau+1}^{-1}(\bar{x}_{\tau+1}(0)) = \begin{cases} h_\tau^{-1}(\bar{x}_\tau(0)) & \text{(a)} \\ h_\tau^{-1}(\bar{x}_\tau(0)) + 1 & \text{(b)} \end{cases} \quad (15)$$

But, by definition, $h_{\tau+1}(x) = h_\tau(x-v)$, so from Eq. (15) one obtains

$$\bar{x}_{\tau+1}(0) = \begin{cases} f_-(\bar{x}_\tau(0)) & \text{(a)} \\ f_+(\bar{x}_\tau(0)) & \text{(b)} \end{cases} \quad (16)$$

where

$$\begin{aligned} f_-(x) &= h_\tau(h_\tau^{-1}(x) - v), \\ f_+(x) &= h_\tau(h_\tau^{-1}(x) - v + 1). \end{aligned} \quad (17)$$

The functions f_- and f_+ inherit some of the properties of h . In particular, f_- and f_+ are continuous and increasing. In the interval $[x^* - a, x^* + a]$ we have that $f_-(x) < f_+(x)$, because h is increasing, so we just evaluate at the following points:

$$\begin{aligned} f_-(x^* + a) &= h_\tau(h_\tau^{-1}(x^* + a) - v) \\ &= h_\tau(i + 1 - \alpha - v), \\ f_+(x^* - a) &= h_\tau(h_\tau^{-1}(x^* - a) - v + 1) \\ &= h_\tau(i - \alpha - v + 1), \end{aligned}$$

where we have made use of Eqs. (14). Thus we have the periodicity condition

$$f_-(x^* + a) = f_+(x^* - a). \quad (18)$$

Next we find when f_- and f_+ reach the extrema of the interval $[x^* - a, x^* + a]$. To this end we determine c_\pm such that $f_\pm(c_\pm) = x^* \pm a$. So we solve

$$\begin{aligned} \begin{cases} f_-(c_-) = h_\tau(h_\tau^{-1}(c_-) - v) = x^* - a \\ f_+(c_+) = h_\tau(h_\tau^{-1}(c_+) - v + 1) = x^* + a \end{cases} \\ \Rightarrow \begin{cases} h_\tau^{-1}(c_-) - v = h_\tau^{-1}(x^* - a) = i - \alpha \\ h_\tau^{-1}(c_+) - v + 1 = h_\tau^{-1}(x^* + a) = i + 1 - \alpha, \end{cases} \end{aligned}$$

whence $h_\tau^{-1}(c_-) = h_\tau^{-1}(c_+)$, and since h is monotonic, we have that $c_- = c_+ = c$.

Therefore, the map Φ giving the dynamics of the central site (12) is given by

$$\Phi(x) = \begin{cases} f_+(x) & \text{if } x^* - a \leq x \leq c \\ f_-(x) & \text{if } x^* + a \geq x > c. \end{cases} \quad (19)$$

From the above properties of f_- and f_+ , it follows that the auxiliary map Φ is a homeomorphism of the circle (see Fig. 8).

A natural binary symbolic dynamics for Φ is introduced by assigning the symbols 0 and 1 whenever the branch f_- or f_+ , respectively, is used in Eq. (15). These symbols correspond to the central site $x(i)$ remaining unchanged, or being replaced by the new site $x(i+1)$, respectively.

Every time a 1 is encountered, the front advances by roughly one site. So the density of 1's in the sequence gives an approximation to the velocity, which becomes exact in the limit $t \rightarrow \infty$. In terms of the circle map, the proportion of 1's in the sequence corresponds to its rotation number ρ :




FIG. 9. Onset of intermittent regime in the auxiliary map, corresponding to the development of a steplike traveling front. For the parameter values and the front shape please refer to Fig. 4(c). The intermittency is the precursor of a pair of period-7 orbits.

$$\rho(\varepsilon) = v(\varepsilon) = \lim_{t \rightarrow \infty} \frac{1}{t} \sum_{i=1}^t s_i, \quad (20)$$

where s_i is the i th term in the symbolic sequence. We have stressed the ε dependence of ρ , since for a fixed local map, Φ depends on ε , and so does its rotation number. Because all sites $\bar{x}(j)$ belong to the same front, the site interchanges all occur at the same time, and therefore the rotation number of any Φ_i is the same as the one for Φ .

The representation of the motion of a front as a circle map implies the likelihood of mode locking for rational velocities, corresponding to Arnold tongues in parameter space, and it affords a simple explanation of the various dynamical phenomena described in the previous sections.

The appearance of a q -period tongue as ε is varied through some critical value ε_* corresponds to a fold bifurcation of Φ^q . Generically, a pair of period- q orbits is created at $\varepsilon = \varepsilon_*$. Thus the orbits of Φ will undergo intermittency in the region of the period- q orbit for ε_n close to ε_* . The intermittency will manifest itself in the graph of Φ as shown by the darkly shaded areas of the orbit web in Fig. 9.

Moreover, the periodic orbit will form toward the center of the dark bands and the corresponding front shape will “flatten” at the heights taken by the periodic points because of the time spent in their neighborhood by the orbits of Φ for $\varepsilon_n \approx \varepsilon_*$. It then follows that the approximating fronts will form steps for the periodic front with the periodic points close to their center points, and independently from the parametric path chosen to approach the boundary point (see Fig. 5).

In Fig. 10 we plot the main mode-locking regions in parameter space (Arnold tongues), corresponding to $v = p/q$ with small q . Here the local map is given by $f(x) = \tanh(x/v)$, while the parameters vary within the unit square: $(\varepsilon, v) \in [0, 1]^2$. We believe that mode locking is a common phenomenon in front propagation in CML's, because the




FIG. 10. Principal Arnold's tongues of the traveling front velocity in the one-way CML with the hyperbolic tangent local map $f(x) = \tanh(x/\nu)$ in the $(\varepsilon, \nu) \in [0, 1]^2$ parameter space. The right hand side scale gives the width σ^2 of the corresponding traveling front for fixed $\varepsilon = 1/2$.

nonlinearity of the local map induces nonlinearity in the auxiliary map [9,10], and mode locking is generic for such maps. However, this phenomenon often takes place on very small parametric scales, since the width of the tongues decreases sharply with increasing ν (Fig. 10). This explains why this phenomenon has not been widely reported (with the notable exception of the large $\nu=0$ region, corresponding to the well-known propagation failure in the anticontinuum limit [18]).

In the continuum limit (see Fig. 10), the stability of the attractors x_{\pm}^* becomes weaker, causing the front to broaden. In Fig. 11 we plotted the auxiliary maps Φ_i corresponding to $\nu = 100/101 \approx 1$ for the one-way CML with local map $f(x) = \tanh(x/\nu)$. This figure should be compared with Fig. 6, corresponding to a narrower front. The domain of each Φ_i is




FIG. 11. Auxiliary maps $\Phi_i(x)$ for the reduced dynamics of the traveling front near the continuum limit. The CML was taken to be one way with local map $f(x) = \tanh(x/\nu)$, $\nu = 100/101$, and coupling strength $\varepsilon = 0.45$.

now smaller, since the interval $I = [x_-^*, x_+^*]$ has to be shared between a larger number of sites. As a consequence, the nonlinearity of each Φ is reduced (note that the auxiliary maps in Fig. 11 are almost linear) and with it the size of the tongues. Thus, the larger the width σ^2 of the traveling front, the thinner the mode-locking tongue (see the right hand side scale in Fig. 10).

ACKNOWLEDGMENTS

R.C.G. acknowledges DGAPA-UNAM (México) for financial support during the preparation of this paper. This work was partially supported by EPSRC GR/K17026.

- [1] K. Kaneko, Prog. Theor. Phys. **72**, 480 (1984); I. Waller and K. Kapral, Phys. Rev. A **30**, 2047 (1984); J. Crutchfield, Physica D **10**, 229 (1984).
- [2] C. Beck, Phys. Rev. E **49**, 3641 (1994); K. Kaneko, Physica D **37**, 60 (1989).
- [3] T. Yanagita and K. Kaneko, Phys. Lett. A **175**, 415 (1993).
- [4] F.H. Willeboordse and K. Kaneko, Physica D **1995**, 101 (1995).
- [5] R. Kapral, R. Livi, G.-L. Oppo, and A. Politi, Phys. Rev. E **49**, 2009 (1994).
- [6] M.P. Hassell, O. Miramontes, P. Rohani, and R.M. May, J. Animal Ecol. **64**, 662 (1995); R.V. Solé and J. Bascompte, J. Theor. Biol. **175**, 139 (1995).
- [7] K. Kaneko, Phys. Rev. Lett. **69**, 905 (1992); Physica D **68**, 299 (1993).
- [8] R. Carretero-González, Ph.D. thesis, Queen Mary and Westfield College, University of London, 1997, <http://www.math.sfu.ca/~nyrcarrete/abstracts.html>
- [9] R. Carretero-González, D.K. Arrowsmith, and F. Vivaldi, Physica D **103**, 381 (1997).
- [10] R. Carretero-González, Int. J. Bifurcation Chaos Appl. Sci. Eng. **7**, 2745 (1997).
- [11] R. Coutinho and B. Fernandez, Nonlinearity **11**, 1407 (1988).
- [12] L.M. Floria and J. Mazo, Adv. Phys. **45**, 505 (1996).
- [13] M. Basler, W. Krech, and K.Y. Platov, Phys. Rev. B **55**, 1114 (1997).
- [14] J. Kosek, I. Schreiber, and M. Marek, Philos. Trans. R. Soc. London, Ser. A **347**, 643 (1994).
- [15] P.C. Bressloff, S. Coombes, and B. deSouza, Phys. Rev. Lett. **79**, 2791 (1997); R. Kuske and T. Erneux, Euro. J. Appl. Math. **8**, 389 (1997).
- [16] P.M. Gade and R.E. Amritkar, Phys. Rev. E **47**, 143 (1993); Q. Zhilin, H. Gang, M. Benkun, and T. Gang, *ibid.* **50**, 163 (1994).
- [17] S.N. Chow and J. Mallet-Paret, IEEE Trans. Circuits Syst., I: Fundam. Theory Appl. **42**, 746 (1995); S.N. Chow, J. Mallet-Paret, and E.V. Vleck, Int. J. Bifurcation Chaos Appl. Sci. Eng. **6**, 1605 (1996).
- [18] S. Aubry and G. Abramovici, Physica D **43**, 199 (1990); R.S. MacKay and J.-A. Sepulchre, *ibid.* **82**, 243 (1995).