

**FINAL Review**

1. Write an explicit integral for the volume  $V$  of the solid obtained by rotating the region bounded by the given curves about the specified line.

$$y = \frac{4}{9}x^2, y = \frac{13}{9} - x^2; \text{ about the } x\text{-axis}$$

2. Use the method of cylindrical shells to write an explicit integral for the volume  $V$  generated by rotating the region bounded by the given curves about the  $y$ -axis.

$$y = 12e^{-x^2}, y = 0, x = 0, x = 1$$

Sketch the region and a typical shell.

3. Evaluate the integral. (Use  $C$  for the constant of integration. Assume  $m \neq 0$ .)

$$\int t \sinh(mt) dt$$

4. Evaluate the integral. (Use  $C$  for the constant of integration.)

$$\int \sin^2(\pi x) \cos^5(\pi x) dx$$

5. Evaluate the integral. (Remember to use absolute values where appropriate. Use  $C$  for the constant of integration.)

$$\int \frac{3x^2 - 19x + 46}{(2x + 1)(x - 2)^2} dx$$

6. Evaluate the integral. (Use  $C$  for the constant of integration.)

$$\int x^3 \sqrt{16 - x^2} dx$$

7. If the infinite curve  $y = e^{-3x}, x \geq 0$  is rotated about the  $x$ -axis, find the surface area of the resulting surface.

8. The air in a room with volume  $180 \text{ m}^3$  contains  $0.15\%$  carbon dioxide initially. Fresher air with only  $0.05\%$  carbon dioxide flows into the room at a rate of  $2 \text{ m}^3/\text{min}$  and the mixed air flows out at the same rate.

Find the percentage of carbon dioxide in the room as a function of time  $t$  (in minutes).

What happens with the percentage of carbon dioxide in the room in the long run?

9. Write TWO explicit integrals, one in  $x$  and one in  $y$ , for the length of the curve  $y = \cos(x)$  for  $0 \leq x \leq \pi$ .  
DRAW A SKETCH!

10. Solve the initial value problem:

$$xy' = y + 3x^2 \sin x, y(\pi) = 0$$

11. Solve the initial value problem:

$$xy' = y^2, y(1) = 1$$

12. Find an equation ( $y = \dots$ ) of the tangent to the curve at the given point.

$$x = \cos t + \cos 2t, y = \sin t + \sin 2t, (x, y) = (-1, 1)$$

13. Find the Taylor series polynomial of order 3 for the function  $y = \sinh(x)$  anchored at  $x_0 = 3$ .

14. Let  $r = f(\theta) = 4 \sin(\theta)$

(A) Sketch the graph of  $r = f(\theta)$  for  $0 \leq \theta \leq 2\pi$  in CARTESIAN coordinates and identify ALL minima and maxima.



(B) Using (A) sketch the graph of  $r = f(\theta)$  in POLAR.

(C)-(D): Use the fact that the slope in parametric is:  $m = \frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}}$ .

(C) Find all  $(r, \theta)$  points where the curve as a HORIZONTAL tangent.

(D) Find all  $(r, \theta)$  points where the curve as a VERTICAL tangent.

(E) By transforming coordinates from polar  $\rightarrow$  Cartesian, find a CARTESIAN equation for this curve.



15. Determine the convergence of the following integral. If convergent, compute its value.

$$\int_0^3 \frac{2}{3-x} dx$$

16. Determine the convergence of the following integral. If convergent, compute its value.

$$\int_0^3 \frac{2}{(3-x)^2} dx$$

17. Determine the convergence of the following integral. If convergent, compute its value.

$$\int_4^\infty \frac{2}{(3-x)} dx$$

18. Find the radius of convergence and interval of convergence of the series.

(a)  $\sum_{n=0}^{\infty} 4^n \frac{(x-2)^n}{n!}$

(b)  $\sum_{n=0}^{\infty} (-1)^{n+1} (n^2 + 1) \frac{(x+5)^n}{3^n}$