

Mode locking of a driven Bose-Einstein condensate

Alexandru I. Nicolin*

Niels Bohr Institute, Blegdamsvej 17, Copenhagen Ø, DK-2100, Denmark

and Theoretical Division and Center for Nonlinear Studies, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA

Mogens H. Jensen†

Niels Bohr Institute, Blegdamsvej 17, Copenhagen Ø, DK-2100, Denmark

R. Carretero-González‡

Nonlinear Dynamical Systems Group, Department of Mathematics and Statistics, and Computational Science Research Center,

San Diego State University, San Diego, California 92182-7720, USA

(Received 21 September 2006; revised manuscript received 6 November 2006; published 13 March 2007)

We consider the dynamics of a driven Bose-Einstein condensate with positive scattering length. Employing an accustomed variational treatment we show that when the scattering length is time modulated as $a\{1 + \epsilon \sin[\omega(t)t]\}$, where $\omega(t)$ increases linearly in time, i.e., $\omega(t) = \gamma t$, the response frequency of the condensate locks to the eigenfrequency for small values of ϵ and γ . A simple analytical model is presented which explains this phenomenon by mapping it to an auto-resonance, i.e., close to resonance the reduced equations describing the collective behavior of the condensate are equivalent to those of a virtual particle trapped in a finite-depth energy minimum of an effective potential.

DOI: [10.1103/PhysRevE.75.036208](https://doi.org/10.1103/PhysRevE.75.036208)

PACS number(s): 05.45.Xt, 03.75.Nt

I. INTRODUCTION

Ever since the experimental attainment of Bose-Einstein condensates (BECs) back in 1995 [1] there has been a surge of research on this topic. The reason for this scientific effervescence is to be found in the almost unprecedented experimental maneuverability of these ultracold gases which engrossed scientists from many distinct fields such as nonlinear dynamics, quantum and nonlinear optics, nuclear and condensed matter physics, to name just a few. Equally appealing are the theoretical insights into the dynamics of Bose-condensed gases obtained through the so-called Gross-Pitaevskii equation (GPE) [2], a cubic Schrödinger equation where the nonlinearity accounts at mean-field level for the interatomic interactions close to absolute zero temperature.

Among the most notable results on the nonlinear side of BECs are the theoretical prediction and the subsequent experimental realization of distinct soliton classes and detailed accounts of their underlying dynamics, see Refs. [3–7] and references therein for some of the recent developments, and the nonlinear infringement of Bloch's periodicity condition. The latter implies that a condensate loaded into the so-called optical lattice, i.e., a periodic potential generated by two counterpropagating laser beams, can have a periodic spatial profile with a period different than that of the underlying lattice [8]. Other works include investigations into the parametric resonances exhibited by a Bose-condensed gas whose scattering length is time modulated on a constant frequency close to the eigenfrequency of the system (see Refs. [9–11], and references therein).

It is the purpose of this paper to investigate the nonlinear dynamics of a repulsive Bose-condensed gas whose scattering length is time-modulated on a linearly increasing frequency, i.e., a scattering length of the type $a\{1 + \epsilon \sin[\omega(t)t]\}$, where $\omega(t) = \gamma t$. While cubic Schrödinger equations have been scrutinized in nonlinear optics over the past few decades, it is, however, only in BECs that one can take advantage of the so-called Feshbach resonances [12] to control the frequency on which the cubic term (i.e., the scattering length) is modulated.

Employing a habitual variational treatment [14] we reduce the dynamics of a three-dimensional fully-symmetric condensate to only one ordinary differential equation (ODE). Our main finding is that, for small values of ϵ , the condensate mode locks to the eigenfrequency, a peculiarity that disappears for high values of ϵ . As shown in Sec. IV, the condensate is collectively described by a single reduced ODE on the condensates' width that emulates the scenario of a particle trapped in the finite-depth energy minimum of an effective potential well.

The paper is structured as follows. Section II is dedicated to the GPE and to the variational method that simplifies the condensate dynamics to an ODE. Section III gives numerical results on frequency locking, while Sec. IV puts forward a simple analytical model that explains the locking through the so-called autoresonance. Section V gathers our conclusions.

II. THE GROSS-PITAEVSKII EQUATION

The dynamics of Bose-condensed gases close to absolute zero temperature is accurately described by the Gross-Pitaevskii equation. In three spatial dimensions it reads

$$i\hbar \frac{\partial \psi}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \psi + V(r)\psi + U|\psi|^2\psi, \quad (1)$$

where $V(r)$ is the trapping potential, here taken as

*Electronic address: nicolin@nbi.dk

†Electronic address: mhjensen@nbi.dk

‡URL: <http://www.rohan.sdsu.edu/~rcarrete/>

$$V(r) = \frac{m}{2}\lambda^2 r^2, \quad (2)$$

where $f = \lambda/2\pi$ is the trap frequency (in Hz) that fixes the strength of the magnetic trap. For simplicity we restrict our analysis to symmetric potentials. The coefficient of the cubic term is $U = 4\pi\hbar^2 a/m$, where a is the two-body scattering length and m the atomic mass.

Due to the intricate nature of the GPE we shall further simplify the problem by restricting ψ to an amenable family of trial functions and study the time evolution of the parameters that define it. This reduces the infinite-dimensional problem of solving Eq. (1) to solving an ODE. A natural choice for the trial function (see Refs. [13,14] for the BEC results, and Refs. [15,16] for similar calculations carried out in nonlinear optics), which actually corresponds to the exact solution in the linear limit ($U=0$), is the three-dimensional Gaussianlike profile

$$\psi(r,t) = A(t) \exp\left[-\frac{r^2}{2w^2} + ir^2\beta\right], \quad (3)$$

where A is the (complex-valued) wave-function at the center of the cloud, w is the width of the condensate while β , the so-called chirp, is the canonical conjugate of w .

After the classical variational recipe (see Refs. [14] for details) the ansatz gives rise to the following equation in the width of the condensate:

$$\frac{d^2}{dt^2}w + \lambda^2 w = \frac{\hbar^2}{m^2} \frac{1}{w^3} + \frac{U}{2\sqrt{2m}} \frac{N}{\pi^{3/2}w^4}. \quad (4)$$

Notice that this equation holds for both time-dependent and time-independent scattering length, i.e., the equation is left unchanged when $U \rightarrow U(t)$. It is worth mentioning that the equation for the width does not depend on the chirp β and that, in turn, the equation for β is driven by the width.

Introducing $P = \sqrt{2/\pi} Na/a_0$, $\tau = \lambda t$ and the rescaled width $v = w/a_0$, where $a_0 = \sqrt{\hbar/m\lambda}$, Eq. (4) reads

$$\frac{d^2}{d\tau^2}v + v = \frac{1}{v^3} + \frac{P}{v^4}. \quad (5)$$

Around the equilibrium point \tilde{v} , defined implicitly by

$$\tilde{v} = \frac{1}{\tilde{v}^3} + \frac{P}{\tilde{v}^4}, \quad (6)$$

the dynamics of the width of the condensate $v = \tilde{v} + \delta$ is given by

$$\frac{d^2}{d\tau^2}\delta + \delta \left(1 + \frac{3}{\tilde{v}^4} + \frac{4P}{\tilde{v}^5}\right) = 0, \quad (7)$$

indicating a period of $T = 2\pi/\omega_P$ where the natural eigenfrequency of the system is

$$\omega_P = \left(1 + \frac{3}{\tilde{v}^4} + \frac{4P}{\tilde{v}^5}\right)^{1/2}. \quad (8)$$

III. MODE LOCKING

Sweeping linearly the frequency of our driving field, which in turn gives a scattering length that goes as $a[1 + \epsilon \sin(\gamma\tau^2)]$, we are faced with the nonlinear ODE

$$\frac{d^2}{d\tau^2}v + v = \frac{1}{v^3} + \frac{P}{v^4}[1 + \epsilon \sin(\gamma\tau^2)]. \quad (9)$$

Equation (9) is solved through an embedded Runge-Kutta method that uses a 4-5 Dormand-Prince pair [17].

Our main finding is that for small values of ϵ and γ (and $\tau < \omega_P/2\gamma$) the width of the condensate shows small amplitude oscillations of frequency equal to the effective frequency of the drive (i.e., $\omega_{\text{osc}} = 2\gamma\tau$). For $\tau > \omega_P/2\gamma$ the width of the condensate shows oscillations of constant amplitude (whose frequency is equal to ω_P , the natural frequency of the condensate) even though the driving field is still present. We call this process ‘‘mode locking’’ [18]. As demonstrated in the next section close to resonance the reduced equations describing the collective behavior of the condensate are equivalent to those of a virtual particle trapped in a finite-depth energy-minimum of an effective potential.

Locking phenomena go a long way back: as early as the 17th century the Dutch physicist Christian Huygens noted that two clocks hanging back-to-back on the wall tend to synchronize their motion. This type of locking is generally present in dissipative systems with competing frequencies. The two frequencies may arise dynamically within the system (as for the two clocks) or through the coupling of an oscillating motion to an external periodic force. In the case of a magnetically trapped BEC (which is usually regarded/ modeled as a nondissipative system) we find that the frequency sweep entailed by the $\sin(\gamma\tau^2)$ term gives rise to a breathing mode whose frequency is equal to the natural frequency of the system.

We illustrate our results on mode-locking by showing the dynamics of the width of the condensate for $P=100$. We purposely chose parameter values that would yield experimental feasible situations. For example, if one considers a magnetic trap with frequency $f \approx 159$ Hz (i.e., $\lambda = 2\pi f = 1000$) for a dilute BEC [13] of approximately $24\,500$ ^{23}Na , the adimensionalization yields a temporal rescaling $\tau = 1000t$ such that, in all our examples, time τ is measured in ms. In Fig. 1 we have plotted the dynamics of the width for $\epsilon=0.01$ and $\gamma=0.05$. This is a typical mode-locking dynamics that includes two distinct regions (see Fig. 1): (i) a transient regime during which the response frequency of the condensate is equal to the effective frequency of the drive (i.e., $\omega_{\text{osc}} = 2\gamma\tau$) and (ii) the mode-locked part when the width of the condensate shows periodic oscillations of constant amplitude. For small γ the critical time τ_c needed to see response on the system’s eigenfrequency may be obtained by writing $\tau = \tau_c + \Delta\tau$ ($|\Delta\tau| \ll 1$) and expanding the argument of the drive $\gamma\tau^2$ yields an effective frequency $\omega_P = 2\gamma\tau_c$, and thus $\tau_c = \omega_P/2\gamma$. Namely, for small γ the transient fades out when the effective frequency of the driving field matches the eigenfrequency of the system. Notice, however, that in Fig. 1 the time needed to stabilize the amplitude is longer as γ is




FIG. 1. (Color online) The dynamics of the width of the condensate for $P=100$, $\epsilon=0.01$, and $\gamma=0.05$. The condensate starts to respond periodically at $\tau_c = \omega_p/2\gamma \approx 22$; the time needed to stabilize the amplitude is, however, slightly longer. This example corresponds to 24 500 ^{23}Na loaded in a magnetic trap with frequency 159 Hz and τ is measured in ms.

somewhat large. Fig. 1 is to be contrasted with Fig. 4 ($\gamma = 0.001$) where the mode-locking takes place exactly at the predicted τ_c .

In Fig. 2 we have plotted the dynamics of the condensate's width for a relatively large value of $\epsilon=1.0$ and $\gamma=0.05$. This is the typical dynamics of the condensate when frequency locking is lost due to the high strength of the driving field. Notice the “bouncing ball” behavior, a peculiarity of the dynamics of the system for high ϵ due to the $1/v^3$ singularity [14].

In Fig. 3 we have plotted the dynamics of the condensate's width for an intermediate value of $\epsilon=0.065$ and $\gamma=0.01$ to illustrate the dynamics outside the linear regime




FIG. 2. (Color online) The dynamics of the width of the condensate for $P=100$, $\epsilon=1.0$, and $\gamma=0.05$. Notice the chaotic response of the condensate and the bouncing ball behavior shown in the inset. Same rescaling as in Fig. 1 so that τ is measured in ms.




FIG. 3. (Color online) The dynamics of the width of the condensate for $P=100$, $\epsilon=0.065$, and $\gamma=0.01$. The condensate starts to respond periodically approximately at $\tau_c = \omega_p/2\gamma \approx 111$ but there are additional superharmonic nonlinear resonances at approximately $\tau_c = n\omega_p/2\gamma$, where $n=2,3,4,\dots$. Same rescaling as in Fig. 1 so that τ is measured in ms.

[19]. The nonlinear regime shows two distinct features: (i) while the observed frequency shows only slight deviations from Eq. (8), the shape of the oscillations shows contributions from the higher harmonics, i.e., strong deviation from the shape of a sine-wave and (ii) in addition to the mode-locking seen approximately at $\tau_c \approx \omega_p/2\gamma$ there is a series of additional super-harmonic nonlinear resonances approximately at $\tau_c = n\omega_p/2\gamma$, where n is an integer larger than one (see Ref. [20]). This latter feature falls outside the simple model put forward in the next section which only accounts for the linear process, i.e., the modelocking at $\tau = \omega_p/2\gamma$.

Finally, in Fig. 4 we have plotted the dynamics of the condensate's width for $\epsilon=0.065$ and small value of $\gamma=0.001$. For this small value of γ the resonances are seen to appear exactly at the predicted $\tau_c = n\omega_p/2\gamma \approx 1115n$ (even for large values of n).

While the reported numerics rely on variational computations we have found the same qualitative behavior in the GPE [21]. Finally, it is important to notice that the frequency locking reported in this paper takes places both for the ground state of the condensate and its excited states. Solitons, for instance, are excited into breather states by the driving field as will be shown elsewhere [21].

IV. ANALYTIC MODEL

Following the recipe of auto-resonance phenomena put forward by Friedland and collaborators (see Ref. [22], and references therein) we recast v as

$$v = \bar{v} + \delta(\tau), \quad (10)$$

where \bar{v} is the equilibrium width [cf. Eq. (6)] and $\delta(\tau)$ measures the deviation from equilibrium. Refraining to the linear regime, i.e., $\delta(\tau) \ll 1$, Eq. (9) reduces to




FIG. 4. (Color online) The dynamics of the width of the condensate for $P=100$, $\epsilon=0.065$, and $\gamma=0.001$. For such a small γ value, the condensate responds periodically precisely at $\tau_c = n\omega_p/2\gamma \approx 1115n$. Same rescaling as in Fig. 1 so that τ is measured in ms.

$$\frac{d^2\delta}{d\tau^2} + \omega_p^2\delta = \frac{\epsilon P}{\tilde{v}^4} \sin(\gamma\tau^2),$$

where we have discarded a term linear in $\epsilon\delta$ for being second order.

We study the dynamics close to resonance and show that the condensate collectively behaves as a particle loaded in a tilted cosinelike effective potential. We show that mode locking is equivalent to the effective dynamics of the particle trapped in an energy minimum of the potential.

Taking $\delta = a(\tau)\sin\varphi(\tau)$ and discarding the second derivative of a with respect to τ , the so-called adiabatic assumption, one has

$$i2\dot{a}\dot{\varphi} + ia\ddot{\varphi} - a\dot{\varphi}^2 + \omega_p^2 a = \frac{\epsilon P}{\tilde{v}^4} \exp(i\gamma\tau^2 - i\varphi),$$

where $\dot{a} = da/d\tau$. Equating real and imaginary parts we obtain

$$a\omega_p^2 - a\dot{\varphi}^2 = \frac{\epsilon P}{\tilde{v}^4} \cos(\gamma\tau^2 - \varphi) \quad (11)$$

for the real part, while the imaginary one gives

$$2\dot{a}\dot{\varphi} + a\ddot{\varphi} = \frac{\epsilon P}{\tilde{v}^4} \sin(\gamma\tau^2 - \varphi). \quad (12)$$

Refraining now to the case close to the resonance, i.e., we limit the analysis to a vicinity of τ_c such that

$$\dot{\varphi}(\tau_c) \approx \omega_p \quad \text{and} \quad \ddot{\varphi}(\tau_c) \approx 0,$$

the previous equations yield

$$\omega_p - \dot{\varphi} = \frac{\epsilon P}{2\omega_p a \tilde{v}^4} \cos(\gamma\tau^2 - \varphi) \quad (13)$$

and

$$\frac{d}{d\tau}(a^2) = \frac{a\epsilon P}{\omega_p \tilde{v}^4} \sin(\gamma\tau^2 - \varphi). \quad (14)$$

Defining the action $I = a^2$ and the phase mismatch $\Phi = \gamma\tau^2 - \varphi$ variables we can recast the previous equations as

$$\dot{\Phi} = 2\gamma\tau - \omega_p + \frac{\epsilon P}{2\omega_p \sqrt{I} \tilde{v}^4} \cos \Phi \quad (15)$$

and

$$\dot{I} = \frac{\sqrt{I}\epsilon P}{\omega_p \tilde{v}^4} \sin \Phi. \quad (16)$$

In order for the condensate to stay mode locked Φ must be close to 0 or π and the right hand side of Eq. (15) should be equal to zero, i.e.,

$$\dot{\Phi}(\tau_c) = 2\gamma\tau_c - \omega_p + \frac{\epsilon P}{2\omega_p \sqrt{I_0} \tilde{v}^4} \cos \tilde{\Phi} = 0,$$

where I_0 is the equilibrium action while $\tilde{\Phi}$ is the equilibrium phase mismatch. Notice that $\dot{\Phi} = 0$ amounts to $\tau_c = \omega_p/2\gamma$, as the last term can be discarded as being numerically small in the small γ limit. The solution of interest is $\tilde{\Phi} = \pi$, for $\tilde{\Phi} = 0$ corresponds to an energy maximum (see below). Setting $I = I_0 + \Delta$ and $\Phi = \tilde{\Phi} + \phi$, where Δ and ϕ are small, the dynamics around the equilibrium is given by the following Hamiltonian system:

$$\dot{\phi} = \Delta S,$$

$$\dot{\Delta} = -A \sin \phi + \frac{2\gamma}{S},$$

where $S = \epsilon P/4\omega_p \tilde{v}^4 I_0^{3/2}$ and $A = \sqrt{I_0} \epsilon P/\omega_p \tilde{v}^4$. The associated Hamilton's function is

$$\mathcal{H}(\Delta, \phi) = \frac{S\Delta^2}{2} + V_1(\phi),$$

where the potential is given by

$$V_1(\phi) = -A \cos \phi - \frac{2\gamma\phi}{S}.$$

If one is to linearize around $\tilde{\Phi} = 0$ the ensuing potential would be

$$V_2(\phi) = A \cos \phi + \frac{2\gamma\phi}{S},$$

which has a maximum for $\phi = 0$ and not a minimum as V_1 has.

The frequency locking reported in the previous section is now transparent: around $\Phi = \pi$ there is an energy minimum corresponding to the system oscillating on its eigenfrequency, while around $\Phi = 0$ there is an energy maximum of no physical interest. Though the linear model discussed above shows an energy minimum independent of the value of ϵ , we conjecture that, in the initial nonlinear problem, the energy minimum is of finite depth and there is a critical

strength of the driving field above which the frequency locking is lost. This process is illustrated in Fig. 2.

V. CONCLUSIONS

In this paper we have shown by means of a variational treatment that modulating the scattering length of a trapped BEC with repulsive interactions as $a[1 + \epsilon \sin(\gamma t^2)]$, leads to the locking of its response frequency to the eigenfrequency for small values of ϵ . Physically, the mode locking amounts to a breathing mode whose frequency is equal to the natural frequency of the system. To the best of our knowledge this is the first paper to analyze mode locking in a BEC context. In order to exhibit the physical mechanism behind it we have restricted ourselves to a variational calculation that captures the main dynamics. To this end we have used a simple analytical model and showed that the equations describing the collective behavior of the condensate are equivalent to those

of a particle trapped in a finite-depth energy minimum of a potential.

Future research should be focused on asymmetric three-dimensional condensates and to analyzing the interplay between the inherent mode-locking processes that take place. Also on the side of future research lies the dynamics of the condensate for negative scattering lengths and the extension of the current observations to multicomponent condensates.

ACKNOWLEDGMENTS

The authors thank Henrik Smith, Christopher J. Pethick, Jan W. Thomsen, Kim Ø. Rasmussen, Mogens Levinsen, and Boris A. Malomed for discussions. R.C.G. acknowledges the support of Grant No. NSF-DMS-0505663. This work was carried out under the auspices of the National Nuclear Security Administration of the U.S. Department of Energy at Los Alamos National Laboratory under Contract No. DE-AC52-06NA25396.

-
- [1] M. H. Anderson, J. R. Ensher, M. R. Matthews, C. E. Weiman, and E. A. Cornell, *Science* **269**, 198 (1995); K. B. Davis, M.-O. Mewes, M. R. Andrews, N. J. van Druten, D. S. Durfee, D. M. Kurn, and W. Ketterle, *Phys. Rev. Lett.* **75**, 3969 (1995); C. C. Bradley, C. A. Sackett, J. J. Tollett, and R. G. Hulet, *ibid.* **75**, 1687 (1995).
- [2] C. J. Pethick and H. Smith, *Bose-Einstein Condensation in Dilute Gases* (Cambridge University Press, Cambridge, 2001).
- [3] J. Denschlag, J. E. Simsarian, D. L. Feder, C. W. Clark, L. A. Collins, J. Cubizolles, L. Deng, E. W. Hagley, K. Helmerson, W. P. Reinhardt, S. L. Rolston, B. I. Schneider, and W. D. Phillips, *Science* **287**, 97 (2000).
- [4] L. Khaykovich, F. Schreck, G. Ferrari, T. Bourdel, J. Cubizolles, L. D. Carr, Y. Castin, and C. Salomon, *Science* **296**, 1290 (2002).
- [5] N. Bilas and N. Pavloff, *Phys. Rev. Lett.* **95**, 130403 (2005).
- [6] S. Sinha, A. Yu. Cherny, D. Kovrizhin, and J. Brand, *Phys. Rev. Lett.* **96**, 030406 (2006).
- [7] S. L. Cornish, S. T. Thompson, and C. E. Wieman, *Phys. Rev. Lett.* **96**, 170401 (2006).
- [8] M. Machholm, A. Nicolin, C. J. Pethick, and H. Smith, *Phys. Rev. A* **69**, 043604 (2004); M. A. Porter and P. Cvitanovic, *Phys. Rev. E* **69**, 047201 (2004); *Chaos* **14**, 739 (2004).
- [9] F. Kh. Abdullaev, E. N. Tsoy, B. A. Malomed, and R. A. Kraenkel, *Phys. Rev. A* **68**, 053606 (2003).
- [10] H. Sakaguchi and B. A. Malomed, *Phys. Rev. E* **70**, 066613 (2004).
- [11] B. Baizakov, G. Filatrella, B. Malomed, and M. Salerno, *Phys. Rev. E* **71**, 036619 (2005).
- [12] S. Inouye, M. R. Andrews, J. Stenger, H.-J. Miesner, D. M. Stamper-Kurn, and W. Ketterle, *Nature (London)* **392**, 151 (1998).
- [13] V. M. Pérez-García, H. Michinel, J. I. Cirac, M. Lewenstein, and P. Zoller, *Phys. Rev. Lett.* **77**, 5320 (1996).
- [14] J. J. G. Ripoll and V. M. Pérez-García, *Phys. Rev. A* **59**, 2220 (1999); B. A. Malomed, *Prog. Opt.* **43**, 71 (2002).
- [15] M. Desaix, D. Anderson, and M. Lisak, *J. Opt. Soc. Am. B* **8**, 2082 (1991).
- [16] E. Caglioti, S. Trillo, S. Wabnitz, B. Crosignani, and P. Di Porto, *J. Opt. Soc. Am. B* **7**, 374 (1990); B. Crosignani, P. Di Porto, and S. Piazzolla, *Pure Appl. Opt.* **1**, 7 (1992); H. Michinel, *ibid.* **4**, 701 (1995).
- [17] E. Hairer, S. P. Norsett, and G. Wanner, *Solving Ordinary Differential Equations. Nonstiff Problems* (Springer-Verlag, Berlin, 1992).
- [18] *Encyclopedia of Nonlinear Science*, edited by Alwyn Scott (Routledge, New York, 2004).
- [19] It is important to stress that while ϵ , is equal to that used in Fig. 1, energy is added to the system due to the small value of γ , therefore the observed deviation from linear case. To see this, one should remember that in the small ϵ , limit the energetic contribution of the driving field goes as $\int_0^\infty \epsilon \sin \gamma t^2 dt = \frac{\epsilon}{\sqrt{\gamma}}$.
- [20] F. K. Abdullaev, J. C. Bronski, and R. M. Galimzyanov, *Photochem. Photobiol.* **184**, 319 (2003); S. Rajendran, P. Muruganandam, and M. Lakshmanan, e-print nlin.ps/0701011.
- [21] A. I. Nicolin, M. H. Jensen, and R. Carretero-González (unpublished).
- [22] J. Fajans, E. Gilson, and L. Friedland, *Phys. Rev. Lett.* **82**, 4444 (1999); *Phys. Plasmas* **6**, 4497 (1999); J. Fajans and L. Friedland, *Am. J. Phys.* **69**, 1096 (2001).